

Climate change and child health: a scoping review and an expanded conceptual framework

Daniel Helldén, Camilla Andersson, Maria Nilsson, Kristie L Ebi, Peter Friberg, Tobias Alfvén



Climate change can have detrimental effects on child health and wellbeing. Despite the imperative for a fuller understanding of how climate change affects child health and wellbeing, a systematic approach and focus solely on children (aged <18 years) has been lacking. In this Scoping Review, we did a literature search on the impacts of climate change on child health from January, 2000, to June, 2019. The included studies explicitly linked an alteration of an exposure to a risk factor for child health to climate change or climate variability. In total, 2970 original articles, reviews, and other documents were identified, of which 371 were analysed. Employing an expanded framework, our analysis showed that the effects of climate change on child health act through direct and indirect pathways, with implications for determinants of child health as well as morbidity and mortality from a range of diseases. This understanding can be further enhanced by using a broader range of research methods, studying overlooked populations and geographical regions, investigating the costs and benefits of mitigation and adaptation for child health, and considering the position of climate change and child health within the UN Sustainable Development Goals. Present and future generations of children bear and will continue to bear an unacceptably high disease burden from climate change.

Introduction

The current impacts and future risks of climate change caused by human activities far surpass those of any other force that has transformed Earth's environment in recent history.¹ Although long-term magnitude and patterns of climate change are uncertain, projections suggest an increase of 2°C or more in the global average temperature could be realised by the end of this century, leading to crucial changes in Earth's geosphere, biosphere, cryosphere, hydrosphere, and atmosphere, with severe implications for human and planetary health.^{2,3} Indeed, climate change will impact the health of children born today throughout their lifetime.⁴

Through its far reaching impact on all parts of society,⁵ climate change will challenge the very essence of children's rights to survival, good health, wellbeing, education, and nutrition⁶ as enshrined by the Convention on the Rights of the Child⁷ and emphasised in the UN Sustainable Development Goals.⁸ Climate change threatens to exaggerate the vulnerabilities of children and other populations at risk and could substantially hamper future progress and possibly even reverse the improvements made in child survival and wellbeing during recent decades.^{6,9}

Despite the clear implications of climate change on child health, Sheffield and Landrigan¹⁰ noted in their 2011 review that there were strikingly few studies examining the possible pathways and mechanisms within this area. The past few years have seen an accelerated momentum for studying the health effects of climate change.^{11,12} Importantly, a comprehensive understanding of how children's health could be affected by climate change, as well as changes in weather patterns and their short and long term impacts, and an identification of strengths, weaknesses, and gaps in the literature to guide future research, policy, and practice are urgently needed. In response to this challenge, we

did a Scoping Review to assess how climate change affects child health and wellbeing.

Methods

Search strategy and selection criteria

Due to the complex nature of the topic and the wide range of studies addressing it, a Scoping Review method was selected^{13,14} based on a predetermined protocol in accordance with established principles (PRISMA Extension for Scoping Reviews)¹⁵ In line with this protocol, a search strategy was developed, and the literature search was done on PubMed, Web of Science, Cochrane Library, Global Health, and WHO regional databases combining two broad blocks of search terms: children ("child", or "infant", or "adolescent") and climate change ("climate change", "global warming", "greenhouse effect", and "climate adaptation" in Medical Subject Heading [MeSH] terms and free text). The decision to not include specific search terms for weather variables (such as temperature) was made because the primary aim of the Scoping Review was to focus on a broad perspective of suggested climate change impacts on child health and not specific variables. Only English language

Key messages

- There is a growing body of research showing that climate change acts through multiple pathways to negatively affect child health and wellbeing
- Despite the imperative for a fuller understanding of how climate change interacts with different determinants to affect child health and wellbeing, gaps persist
- A range of research methods are needed to further understanding of vulnerable subpopulations and geographical regions, and of the costs and benefits of mitigation and adaptation for addressing risks

Lancet Planet Health 2021;
5: e164-75

Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden (D Helldén MD, T Alfvén PhD); Department of Epidemiology and Global Health, Umeå University, Umeå, Sweden (C Andersson MSc, Prof M Nilsson PhD); Department of Global Health, School of Public Health, University of Washington, Seattle, WA, USA (Prof K L Ebi PhD); Swedish Institute for Global Health

Transformation, Royal Swedish Academy of Sciences, Stockholm, Sweden (Prof P Friberg PhD); Department of Public Health and Community Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden (Prof P Friberg); Sachs' Children and Youth Hospital, Stockholm, Sweden (T Alfvén)

Correspondence to: Mr Daniel Helldén, Department of Global Public Health, Karolinska Institutet, Stockholm 17177, Sweden daniel.hellden@ki.se




Figure 1: Study selection process
Documents refer to the material as a whole (original articles, reviews, commentaries, editorials, and grey literature).




Figure 2: Number of original articles published between 2000 and 2019
In 2019, only published articles before June 11 were counted.

studies from Jan 1, 2000, to June 11, 2019, were considered. The detailed search strategy and PRISMA Extension for Scoping Reviews are available in the appendix (pp 2–3).

See Online for appendix

A screening of the retrieved literature was done by DH by reviewing the titles and abstracts in terms of relevance to the research question—that is, the inclusion criteria being that the original article, review, commentary, editorial, or grey literature (collectively referred to as documents henceforth) found were concerned with children (as defined by the Convention on the Rights of the Child⁷ as people younger than 18 years) and a climate-related exposure that could be a risk factor for child

health. In this Scoping Review we assumed the statements concerning the exposure were accurate with respect to whether they were weather, climate, climate variability, or climate change as defined by the Intergovernmental Panel on Climate Change.¹ The relevant documents were considered for full-text review by DH and CA. The documents included for full-text review were assessed for eligibility; ineligible documents did not have a full text available, were not in English, did not concern children, or did not include variables or analyses relevant to climate change as previously defined. To validate the process, MN and TA independently reviewed a random sample of 10% of the documents included for full-text review, and any discrepancies with regards to inclusion were discussed by DH, CA, MN, and TA to reach consensus.

Data abstraction

A first data abstraction chart based on published frameworks on health and climate change^{10,16,17} was developed and tested by DH and CA. This chart included generic document information; contextual aspects of the original articles and reviews such as study setting, study groups, and information on climate change-related exposures and climate-sensitive health outcomes, including heat-waves, air pollution, food and nutrition, infectious diseases, and mental health; and social determinants of health, such as poverty and violence. After inputting data from the included documents, DH and CA applied open coding¹⁸ to revise the categories of the chart and establish key themes across the included material. From this chart, the conceptual framework was created and revised by all authors. Due to the large number of studies included in the Scoping Review and their different approaches, a critical appraisal of the quality of each study was deemed to be beyond the scope of this study. The few original or review articles (n=14) that exhibited severe quality gaps in the form of unspecified methods determined by the full-text review were hence reviewed but not included in the analysis. As a consequence of the large material considered, only the most relevant are referenced throughout the result section below; a list of all included documents can be found in the appendix (p 5).

Role of the funding source

The funder of the study had no role in study design selection, data collection, data analysis, data interpretation, or writing of the report. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

The initial search yielded 2970 documents, of which 749 were duplicates, leaving 2221 documents that were screened for relevance (figure 1). Of these, 1705 were excluded, resulting in 516 documents being considered for a full-text review. From these, 145 were excluded due

to their full text being unavailable, they were not in English, or they did not include either a child or climate change component. The total number of 371 documents were reviewed (202 original articles, 118 reviews, and 51 documents of other types such as commentaries, editorials, and grey literature). There has been a marked increase in the number of original articles per year on the topic, as shown in figure 2. In 2019, only published articles before June 11 were counted.

As described in the table, the largest proportion of single country studies were in high-income countries, with only 12.2% in low-income countries, primarily done in countries in Asia and Africa. Approximately half of the studies included urban and rural settings, and almost all used some kind of quantitative analysis. Only a third of the articles disaggregated the results by sex. The age group(s) used to define a child were heterogeneous and in some instances not explicitly stated, but the majority of studies focused on children aged younger than 5 years.

The effects of climate change on child health

A conceptual overview of how a changing climate could affect child health is presented in figure 3. The framework details how greenhouse gas emissions lead to a changed Earth system with higher temperatures, altered rainfall, a rise in sea level, acidification of oceans, and an increased frequency of rapid and delayed onset extreme weather events. Direct effects of climate change include temperature changes (heatwaves and more rapidly changing temperatures), changing precipitation patterns with increased risk of floods, droughts, and wildfires. More indirect effects include ecosystem disruption, changing vector patterns, air pollution, and aeroallergens. Child health risks due to these effects can be categorised as: a cause of direct harm to children and an increase in the risk of disease. Figure 3 also incorporates how these effects interact with human activity and the concepts of mitigation and adaptation.

Direct effects

The direct effects of climate change in the form of changing weather patterns—for instance, rapid and delayed onset extreme weather events such as storms, heatwaves, and flooding, are associated with childhood morbidity and mortality.¹⁹ However, the underlying evidence of how climate change could directly affect adverse health outcomes is limited.^{10,20}

Temperature changes

Children could be particularly at risk for negative health outcomes from rapid or delayed heatwaves.²¹ Heatwaves increase overall mortality rates in a population.²² Xu and colleagues²³ found inconclusive evidence of a higher risk of child mortality during heatwaves but the evidence was somewhat stronger for a heightened risk to infants than older age groups. Subsequent studies emphasised an increased overall mortality risk from heatwaves in

% (n)	
Income status (single country original articles, n=172)	
Low income	12.2 (21)
Low-middle income	22.1 (38)
Upper-middle income	24.4 (42)
High income	41.3 (71)
Geographical area (single country original articles, n=172)	
North America	19.8 (34)
South America	5.2 (9)
Europe	4.7 (8)
Asia	36.0 (62)
Africa	22.7 (39)
Oceania	11.6 (20)
Study setting (all original articles, n=202)	
Urban	29.7 (60)
Rural	19.3 (39)
Both	51.0 (103)
Study method type (all original articles, n=202)	
Quantitative	92.6 (187)
Qualitative	4.0 (8)
Mixed method	3.5 (7)
Results disaggregated by sex (all original articles, n=202)	
Yes	32.2 (65)
No	67.8 (137)

Table: Characteristics of the original articles included

children, particularly infants,^{24,25} and multiple studies in low-income and middle-income settings^{26–28} quantified the detrimental effect of heatwaves on child mortality rates, with some exceptions to this observed association.²⁹ Direct morbidity from heatwaves include heatstroke, electrolyte imbalance, kidney-associated diseases, and respiratory as well as infectious diseases,^{23,30,31} whereas increases in mean temperature are associated with a higher prevalence of paediatric kidney stones and Kawasaki disease. Emergency department visits by children in cities increase during heatwaves,³² especially among infants,³³ with the magnitude of risk differing between and within countries.³⁴ Asthma might be the most common cause for heatwave-associated respiratory disease.^{35,36} Heatwaves might also result in an increase in unintentional injuries.³⁷ Combining socioeconomic and demographic factors with spatial modelling can forecast at-risk locations for increased heatwave-related mortality and morbidity.³⁸

Increasing temperatures are not the only aspect that harm children. Studies describe greater mortality from cold temperatures among children in Madrid, Spain, and China,^{39,40} whereas a rapid change of temperature and humidity levels can significantly affect respiratory and infectious diseases in children.^{41,42}

Precipitation and floods

Rapid onset storms and heavy precipitation compound the risk of flooding and subsequent direct mortality and




Figure 3: Climate change and child health: an expanded framework
 The dashed lines show where mitigation and adaptation can hinder the effects of climate change on child health and wellbeing. HFMD=Hand, foot, and mouth disease. *Other infectious diseases, from mumps to bacterial meningitis, varicella viruses, and parasitic diseases.

morbidity from drowning and unintentional injuries.^{43,44} For child health, increases in precipitation are associated with overall higher mortality rates⁴⁵ and have long-term effects on child health due to lack of proper nutrition and the spread of communicable diseases. Flooding increases childhood mortality and morbidity^{43,46} through amplified risk of infectious diseases, especially from water and food,^{46,47} malaria, and respiratory infections, as well as its effect on the overall nutrition status of children,⁴⁸ and thus disproportionately affects the most vulnerable.

Droughts and wildfires

A lack of precipitation over time along with many other factors, such as evapotranspiration associated with increased temperatures, can lead to droughts. Droughts generally have a longer onset and duration than other weather and climate events.⁴⁹ Putting a strain on basic water needs for families in rural settings, droughts and drought-associated wildfires also contribute substantially to worsen respiratory health,⁴³ mental health,⁵⁰ under-nutrition,⁵¹ and infectious diseases among children.⁵² Long-term changes in rainfall patterns, most notably lesser annual volume, affect child mortality rates in the Sahel region, Africa.⁵³ Despite potential adaptation strategies, droughts undermine local ecosystems, drive families into poverty, and can force migration.⁴⁹

Indirect effects

Most indirect effects of climate change are multifaceted, acting over short and long timescales. These include ecosystem disruption, changing vector patterns, air

pollution, and aeroallergens. Acting through a myriad of causes linked to climate change, mental health among children are also reviewed in this section.

Changing vector patterns and infectious diseases

The distribution of disease-carrying vectors are expected to be altered by climate change, whereas the transmission and survival of infectious pathogens will be influenced by changing weather patterns.^{54,55} However, identifying specific infectious diseases that are sensitive to climate change remains difficult. Infectious diseases take a disproportionately high toll on children in general and socioeconomically vulnerable children in particular.⁵⁶

The relationship between local or regional weather trends and the geographical range, seasonality, and prevalence of malaria is well understood. The long-term impact of climate change on the distribution of malaria vectors, the parasite itself, and child mortality and morbidity are under investigation.⁵⁷ In Uganda, short-term weather trend changes, plausibly due to climate change, leading to high temperatures and increased flooding, have led to more favourable conditions for malaria transmission, hampering vector control efforts.^{58,59} Similar results have been found in Mali and Zambia. Generalising the net effect of climate change, intensified malaria transmission across sub-Saharan Africa could increase the malaria related under-5 mortality by 10–15% if no measures are taken to mitigate or adapt to climate change.⁵⁷ Using geospatial modelling based on the representative concentration pathways with the highest greenhouse gas emission (Representative Concentration

Pathway [RCP]), Semakula and colleagues⁶⁰ suggested that areas with high malaria prevalence will move towards eastern and southern rural Africa during this century.

Climate variability, particularly average temperature change, is associated with a change of geographical distribution of dengue,⁶¹ with few quantifications of the associations specifically for children, although the burden of disease and deaths caused by the virus is highest among children in rural settings. One study estimated that almost half of the world's population could be at risk for dengue fever by the end of the century.⁶²

Most prevalent in North America and Europe, the *Borrelia burgdorferi* bacteria that causes Lyme disease is transmitted by ticks, with children often being at most risk.⁶³ The lifecycles of the bacteria and tick are dependent on temperature and humidity, among other factors, and rising temperatures with climate change have expanded the geographical areas of transmission of Lyme disease into more northern parts of North America and Europe, and increase the risk of transmission in endemic areas.^{54,64} A recent study projected a 20% increase in the incidence of the disease in the USA by mid-century compared with the current incidence rate under RCP 4.5.⁶⁴

The multifactorial aetiology of diarrhoea has made it difficult to attribute cases to climate change,^{62,65} a systematic review of the association between temperature and all causes of diarrhoea suggested however that climate change could be responsible for a substantial portion of diarrhoea cases.⁶⁶ Specifically, the prevalence of *Escherichia coli*, *Cryptosporidium*, rotaviruses, and parasites (eg, *Giardia* and *Toxoplasma gondii*) increase with higher temperatures across many different climatic zones. Rural areas experiencing lower amounts of rainfall suffer from higher diarrhoea incidence rate among children due to the use of unsafe drinking sources and limited hygiene.^{52,67} Conversely, higher precipitation levels in urban areas, both in low-income and high-income settings might increase diarrhoea incidence due to sub-optimal water and sanitation systems.^{68,69} A warmer and dryer climate under RCP 8.5 could increase the intensity of rainfall in countries such as Botswana, South Africa, Senegal, Rwanda, and Ethiopia possibly leading to changes in patterns of childhood diarrhoea prevalence, underscoring the need for effective water and sanitation systems. Studies from India show that rainfall is associated with an increased prevalence of childhood diarrhoea; extrapolating past changes in climate could mean a 13% increase in childhood diarrhoea incidence rates in the northern states by 2040.⁷⁰ Increasing numbers of heavy rainfall and floods in Cambodia and China show that these events put children at risk for diarrhoea.^{47,71} Evidence from other Asian countries emphasises that temperature is the most important weather variable associated with diarrhoea although this differed between geographical areas.^{72,73}

Climate variability in the form of increased rainfall and humidity correlates with a higher risk of contracting

bacterial and viral lower respiratory tract infections among children.⁷⁴ Studies from Australia and China show that rapid onset extreme weather events such as heatwaves, low temperatures, and heavy precipitation increase the vulnerability of children to a range of pathogens causing such infections.⁷⁵ Climate change will affect patterns of respiratory pathogens in both rural and urban settings.⁷⁴

Hand, foot, and mouth disease (HFMD) is caused by enteroviruses that primarily infect children, and the prevalence of HFMD has increased in southeast Asia.⁷⁶ Examining the association between weather variables and HFMD suggests that the increased incidence of HFMD in China, Vietnam, and Japan is associated with higher average temperatures, precipitation, humidity, and air pollution in urban settings. Zhao and colleagues⁷⁷ have projected that the incidence of HFMD could increase by 3–5% by the end of the century in China, but the main impact would be a change in which urban areas would be prone to HFMD outbreaks. The trend and pattern appear to be true for other southeast Asian countries as well.⁷⁸ A changing climate could also affect the geographical range of other infectious diseases, from mumps to bacterial meningitis, varicella viruses, and parasitic diseases (eg, schistosomiasis).

Air pollution and aeroallergens

The effect of exposure of children to air pollution and aeroallergens, particularly on their respiratory systems, is detrimental. The air pollutants of particular concern with climate change are particulate matter and ozone. Aeroallergens will shift in seasonality and concentrations with climate change.

Particulate matter air pollution is a direct by-product of fossil fuel combustion and will worsen with climate change-related alterations in droughts, fires, and sand storms.^{55,79} Exposure to particulate matter adversely affects lung function in children, with decreased peak expiratory flow and forced expiratory volume, particularly for children with asthma, leading to an increased number of emergency department visits, hospital admissions, and deaths in children.^{80,81} Infant and child mortality reductions due to access to electricity in resource-limited and rural settings might actually be offset by air pollution from biomass and petroleum sources.^{82,83} There could be an association between exposure to particulate pollution during pregnancy and children being small for gestational age, having low birthweight, and later presenting respiratory disorders,^{84,85} although the results across individual studies are not consistent. Perera and colleagues⁸⁶ developed dose response curves between exposure to particulate matter during pregnancy and preterm birth, low birthweight, and incidence of asthma. There seems to be a reinforcing relationship between particulate matter and high temperatures and low humidity leading to a slightly higher risk of mortality and morbidity, but a more complex relationship regarding other climatic variables.

With increasing mean temperatures, constant emissions of precursor chemicals, such as nitrogen dioxide from fossil fuel combustion and biological processes, and no change in cloud cover, ozone concentrations are projected to increase especially in urban settings, but the extent will depend on local climatic variables.⁸⁷ Higher concentrations of ozone might exacerbate respiratory disorders such as asthma, atopic dermatitis, and allergic conjunctivitis. In South Korea, ozone pollution is projected to cause the majority of the burden of disease from climate change in children younger than 15 years during this century, assuming a balanced climate change model (A1B).⁸⁸ Other air pollutants from fossil burning sources such as transportation that can have a clear impact on child health both short term and long term are carbon monoxide, sulfur oxides, and nitrogen dioxide which similarly to ozone can worsen the respiratory function in children particularly for those suffering from asthma or other chronic respiratory diseases.^{80,89}

Climate change and higher carbon dioxide concentrations, through increased mean temperature and greater precipitation, prolong and intensify the pollen season and affect the severity of allergic rhinitis, allergic conjunctivitis, atopic dermatitis, and asthma.^{89,90} Understanding the burden of allergenic diseases due to climate change is difficult due to multifaceted causal pathways.⁹¹ Neumann and colleagues⁹² projected that emergency department visits from asthma caused by aeroallergens could increase by 14% in the USA over the course of this century under RCP 8.5 due to a longer pollen season. Additionally, warmer and more humid weather may increase the risk of fungal spore production, indoor mould growth, and insect and contact allergies.

Mental health

The implications of climate change on mental health of children has been poorly researched, although it has come under increased scrutiny during the last decade.^{93,94} Burke and colleagues⁹³ suggested that climate change in the form of more frequent rapid and delayed onset extreme weather events and disruption of socioeconomic factors increases young people's susceptibility to mental health illness in the form of depression, post-traumatic stress disorder, and anxiety, and that climate change might lead to graver mental health illnesses, particularly among already disadvantaged children. Empirical qualitative studies on the effect of drought^{50,95} and higher temperatures⁹⁶ on adolescent mental health in Australia and the USA show an overall worse mental health with higher levels of emotional and physiological distress. Studies from Sweden,^{97,98} Canada,⁹⁹ and Australia¹⁰⁰ showcase how adolescents coped with the disruptions by focusing on the problem or building meaning around combating climate change. It has also been suggested that the effects of climate change might have long-term behavioural effects.

Increasing the risk of disease

Through direct and indirect mechanisms and over long and short timescales, climate change can affect determinants of general child health and wellbeing. The key determinants found in this Scoping Review are detailed in this section.

Food insecurity and malnutrition

Malnutrition is one of the leading factors contributing to child mortality and morbidity globally.^{54,56} The empirical evidence on how climate change will impact child health through the effect on nutrition is under development but mainly focused on rural areas.¹⁰¹ The drivers of and interactions between undernutrition and overnutrition and climate change are complex.¹⁰²

The majority of studies indicate that potential gains in crop yields from higher average temperatures and carbon dioxide concentrations would be offset by negative effects in the form of heatwaves, lower water availability, changes in the dynamics of crop diseases, and lower nutritional value of crops (for instance lower concentration of iron in foods).^{54,103,104} Extreme weather events such as flooding have long-term effects on the nutrition status of children, with the most disadvantaged children being at greatest risk.^{48,105,106} Studies in west Africa, Ethiopia, Kenya, and around the Lake Victoria Basin, where the crop yield and other agricultural variables were compared with undernutrition in children, showed that decreases in such variables due to droughts lead to a higher probability of child undernutrition. A number of studies modelled changes in temperature and precipitation due to climate change based on RCP 8.5 over the coming decades, and projected that sub-Saharan Africa could face increased rates of stunting of up to 20% depending on rate of socioeconomic development^{101,107,108} but that this would vary substantially within countries. Crop diversification has been identified as a possible adaptation measure in combating child undernutrition.¹⁰⁹ The rapid economic growth of Asian countries has led to lasting undernutrition in rural areas and overnutrition in cities.¹¹⁰ Lastly, studies have emphasised that there are macro and micro economic, environmental, technological, cultural, and societal determinants that influence the drivers of child undernutrition due to climate change,¹¹¹ and that the economic impact might be negligible.¹¹²

Patterns of nutritional intake by children and the built environment interact to facilitate unhealthy diets and decreased levels of activity, leading to increased risk of overnutrition.¹¹³ Similar to undernutrition, the risk factors associated with overnutrition have been found to be most prevalent among socioeconomically vulnerable children in urban settings.¹¹³ Indeed, overnutrition, undernutrition, and climate change seem to have a common set of causes requiring a systematic approach to finding and implementing solutions.¹⁰²

Socioeconomic distress, education, migration, and toxicant exposure

Not many studies directly examine the more distant determinants of child health and their relation to climate change. It is clear, however, that the health effects of climate change will magnify already existing child health inequities.

Climate change could force families into poverty through loss of livelihood, additional costs from property damage, and health costs.⁵⁴ Particularly vulnerable are populations already living in poverty or near poverty in rural geographical areas that will probably be affected most from climate change such as low-elevation coastal zones due to sea level rise, less-favoured agriculture areas, or the Arctic region. The wealth of individuals and families will be important factors when adapting to climate change.¹¹⁴ The impact of climate change-related extreme weather events on child poverty rates seems to be indifferent to the quality of governance structures.¹¹⁵ Climate change affects the broader societal context, with studies describing how disruptions such as storms in Bangladesh have increased the risk of early child marriage as a way of coping with economic loss¹¹⁶ and that gender violence weakens the resilience of communities.¹¹⁷ Another vulnerable subgroup of children are those who lack a social security net.^{118,119} The many and different social, economic, and environmental determinants that are affected by climate change often amplify the health impact on children particularly in low-income settings.¹²⁰ At the same time, children are themselves agents of change and an integral part of the resilience of families and communities.

Climate variability, including higher temperatures during prenatal and early childhood years, can lead to significantly less educational attainment, but the relationship between rainfall and level of education differs across regions and rural and urban settings.¹²¹ High ambient air temperature during schooling leads to negative effects on the wellbeing of children,¹²² whereas females experiencing droughts during childhood have lower educational attainment.⁵¹ In turn, low education has often been cited as one reason for low socioeconomic status that leads to disproportionately higher risk from the effects of climate change. The potential of schools in addressing health impacts from climate change is largely unknown.

Few articles investigated migration as a result of climate change and its impact on child health, but it is expected that losses of sustainable water sources due to sea level rise and flooding could be one of the key drivers of migration.¹²³ Examining the migration pattern in response to natural disasters in Bangladesh, Penning-Rowsell and colleagues¹²⁴ concluded that migration occurred either for fear of safety or for economic reasons, but that permanent migration was only seen as a last resort.

Due to the increased frequency of extreme weather events and overall changes in temperature and

precipitation from climate change, it is probable that children's exposure to toxicants such as organic pollutants, pesticides, metals, and other toxicants could increase.⁵⁵ Exposures to aforementioned toxicants can not only lead to acute poisonings, but also to long-term adverse health effects such as endocrinological or neurological dysfunctions.¹²⁰

In-utero exposure

The effects of climate change on pregnant mothers are beginning to be understood, with studies focusing on the intersection between maternal and child health with regards to climate change.¹²⁵

Extreme temperature changes during pregnancy have been associated with an increased risk of preterm birth, but the association varies across studies,^{126,127} and does not appear to be significant in some high-income contexts.¹²⁸ Water scarcity due to climate variability has also been identified as increasing the risk of preterm birth.¹²⁹ Extreme temperatures, low amounts of precipitation, and high levels of air pollution have led to lower birthweight and affected length at birth, which has implications for the health of the neonate and development of the child.^{84,128,129} The magnitude and association between climatic variables and birth status vary, but climate change could lead to lower average birthweights in sub-Saharan Africa over the coming decades.¹³⁰ Some studies suggest that higher average temperatures increased the prevalence of birth defects.¹³¹ There are also indications for a connection between extreme temperatures and miscarriages and stillbirths,^{132,133} whereas higher water salinity at the end of the pregnancy from rises in sea level or flooding points to an increase in the risk of neonatal death.¹³⁴ Rapidly changing and extreme temperatures during pregnancy was linked to childhood pneumonia,^{135,136} air pollution, and childhood atopic eczema,¹³⁷ whereas mental health illness during pregnancy following rapid onset extreme weather events impacted infant behaviour¹³⁸ and child height. There are also some indications that increased in-utero exposure to particular matter and other air pollutants could have physical, cognitive, and behavioural effects on the child later on in life.⁸⁶ Early evidence showed that high temperature exposure in utero led to a small reduction in adult wages in the USA;¹³⁹ however, the exact implications for other outcomes remain unclear.

Discussion

Climate change and child health is a new field of inquiry, with the first review dedicated to the subject published in 2003¹⁴⁰ and subsequently advanced by the American Academy of Paediatrics in 2007,¹⁴¹ as well as Ebi and Paulson in 2007 and 2010.^{63,142} Our review assesses the current evidence on how climate change can affect child health, synthesising a vast body of literature, and presents an expanded framework that can be used for understanding and positioning climate change and child

health within the sustainable development discourse. Of the 202 original articles included, almost all were published during the last 5 years. Albeit having a relatively equal geographical distribution, most of the studies were confined to upper-middle-income or high-income countries, consistent with previous evidence summaries.^{10,16,54,55,63,142–144}

The prevailing inequalities, between and within countries, largely determine how climate change impacts children. Disadvantaged children suffer from a disproportionately high and truly unjust health burden from climate change.¹⁴⁵ Further, the impact of climate change seems to follow an urban and rural divide, although not consistently. This pattern is probably due to the differences in the environment,¹⁴⁶ access to health care, and socio-economic factors.¹⁴⁷ The difficulty of deciphering weather variability due to climate change and the attributable health impact was acknowledged by almost all studies. Other factors might cause a certain outcome, and there remains a great level of uncertainty when it comes to measures of climate change health impacts on children.

Previously identified research gaps within the area of climate change and child health have not been addressed to satisfaction. First, the large number of articles found masks a striking lack of focus on climate change and child health; many studies only include children as a subpopulation of analysis. Second, although there exists research from low-income settings, most studies were still done in high-income and upper-middle-income countries and focused on particular countries. Third, there was little evidence of the beneficial or harmful effect of social, political, and commercial determinants on climate change impacts on the physical and mental health of children and different scenario projections taking these and different climate change scenarios into account. Fourth, few articles disaggregated the analysis by sex and age group, limiting the generalisability and usability of results. Fifth, the health outcomes that are the basis of the assessment of climate change impact on children vary to such a large degree that it is almost impossible to compare results across studies. Last, there is a clear lack of qualitative research and longitudinal studies, restricting a fuller understanding of how climate change affects child health and wellbeing.

Several potential areas for future investigation were found. The use of temporo-spatial modelling, providing detailed estimates of child mortality,¹⁴⁸ could untangle complex relationships between climate change, weather patterns, and health impacts. An unexplored area comprises the co-benefits of climate change mitigation and reduced ill health for children, concretely exemplified by the linkage between fossil fuel use both at a household and societal level and subsequent air pollution.⁸³ There is also a great need for studying adaptation, with vital societal functions such as access to safe and quality water and sanitation services being key for developing resilience against climate driven infectious disease

outbreaks. Situating climate change effects within the framework of the UN Sustainable Development Goals further enables the holistic exploration of interlinkages between climate change, child health, and other societal sectors.¹⁴⁹ The research gaps found should not be an excuse for inaction; our findings show that there is clear impetus for policy makers to strengthen mitigation and adaptation efforts.

A clear strength of this Scoping Review is the bringing together of a wide variety of studies using a systematised search strategy providing a broad overview of the area. When trying to synthesise such a complex topic there are certain limitations and potential for biases. First, we restricted the search in terms of time, language, and few search terms, which inherently have led to excluding some studies that did not include the search terms as MeSH terms or free text. As with all literature searches, there is a risk of publication bias, which, due to the large number of studies indicating causal relationships found, is deemed likely. Second, we did not publish the protocol of the study due to time limitations and only one investigator (DH) did the screening process and, although validated procedures were in place, the inclusion process could be limited by selection bias. However, the data extraction and quality assessment were done by two investigators (DH and CA) in cooperation to ensure consistency and we explicitly state which documents we refer to in this review and make all included documents available in the appendix (p 5). Third, we do not provide any appraisal of the level of uncertainty for each association described, if the authors accurately described their association as weather, climate variability, or climate change or a definitive conclusion on the potential heightened risk or similar measurement; hence, the actual impact from climate change and the described aspects cannot be asserted within a particular population. In summary, the strengths and limitations of this review mirror the challenges and opportunities within the field of climate change and child health.

The effects of climate change on child health travel through many different pathways, vary substantially across geographical locations, and are heavily influenced by broader socioeconomic contexts. By using a diverse range of novel research methods, studying overlooked populations and geographical regions, and investigating the benefits of mitigation and adaptation to child health, crucial knowledge gaps can be filled. Rejuvenating innovative research efforts across disciplines is crucial to fill these gaps, where situating climate change and child health within the UN Sustainable Development Goals will be key. Still, present and future generations of children, particularly the already vulnerable, bear and will continue to bear an unacceptably high disease burden from climate change.

Contributors

TA and MN generated the study idea. DH, CA, TA, and MN designed the analysis and methods. DH and CA reviewed the literature. DH and CA

did the analysis, and DH wrote the first draft of the manuscript, to which all authors contributed. KLE and PF participated in the analysis and provided critical scientific input on the manuscript.

Declaration of interests

DH, CA, MN, and TA report grants from Formas (FR-2018/0010) during the conduct of the study. All other authors declare no competing interests.

Data sharing

All data are available in the paper and appendix.

Acknowledgments

We would like to thank Ylva Bergman at the Karolinska Institutet library for assistance with developing the search strategy and Göran Tomson for suggestions to improve the manuscript.

References

- Intergovernmental Panel on Climate Change. Special report: global warming of 1.5°C. Geneva: World Meteorological Organization, 2018.
- Collins M, Knutti R, Arblaste J, et al. Long-term climate change: projections, commitments and irreversibility. In: Stocker TF, Qin D, Plattner GK, et al, eds. Climate Change 2013—the physical science basis: working group I contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Geneva: World Meteorological Organization, 2013.
- Whitmee S, Haines A, Beyrer C, et al. Safeguarding human health in the Anthropocene epoch: report of The Rockefeller Foundation-Lancet Commission on planetary health. *Lancet* 2015; **386**: 1973–2028.
- Watts N, Amann M, Arnell N, et al. The 2019 report of The Lancet Countdown on health and climate change: ensuring that the health of a child born today is not defined by a changing climate. *Lancet* 2019; **394**: 1836–78.
- Intergovernmental Panel on Climate Change. AR5 climate change 2014: impacts, adaptation, and vulnerability. Cambridge, UK and New York, NY, USA: Cambridge University Press, 2014.
- UNICEF. Unless we act now: the impact of climate change on children. 2015. https://www.unicef.org/publications/files/Unless_we_act_now_The_impact_of_climate_change_on_children.pdf (accessed Nov 20, 2019).
- UN. Convention on the Rights of the Child. 1989. <https://www.ohchr.org/Documents/ProfessionalInterest/crc.pdf> (accessed Nov 20, 2019).
- UN General Assembly. Transforming our world: the 2030 agenda for sustainable development. 2015. https://www.un.org/en/development/desa/population/migration/generalassembly/docs/globalcompact/A_RES_70_L_E.pdf (accessed Nov 20, 2019).
- WHO. United Nations Environment Programme. Healthy environments for healthy children: key messages for action. Geneva: World Health Organization and United Nations Environment Programme, 2010.
- Sheffield PE, Landrigan PJ. Global climate change and children's health: threats and strategies for prevention. *Environ Health Perspect* 2011; **119**: 291–98.
- Verner G, Schütte S, Knop J, Sankoh O, Sauerborn R. Health in climate change research from 1990 to 2014: positive trend, but still underperforming. *Glob Health Action* 2016; **9**: 30723.
- Watts N, Amann M, Ayeb-Karlsson S, et al. The Lancet Countdown on health and climate change: from 25 years of inaction to a global transformation for public health. *Lancet* 2018; **391**: 581–630.
- Pham MT, Rajić A, Greig JD, Sargeant JM, Papadopoulos A, McEwen SA. A scoping review of scoping reviews: advancing the approach and enhancing the consistency. *Res Synth Methods* 2014; **5**: 371–85.
- Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* 2018; **18**: 143.
- Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018; **169**: 467–73.
- Philipsborn RP, Chan K. Climate change and global child health. *Pediatrics* 2018; **141**: e20173774.
- Watts N, Amann M, Arnell N, et al. The 2018 report of the Lancet Countdown on health and climate change: shaping the health of nations for centuries to come. *Lancet* 2018; **392**: 2479–514.
- Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008; **8**: 45.
- Garcia DM, Sheehan MC. Extreme weather-driven disasters and children's health. *Int J Health Serv* 2016; **46**: 79–105.
- Amegah AK, Rezza G, Jaakkola JJK. Temperature-related morbidity and mortality in sub-Saharan Africa: a systematic review of the empirical evidence. *Environ Int* 2016; **91**: 133–49.
- O'Neill MS, Ebi KL. Temperature extremes and health: impacts of climate variability and change in the United States. *J Occup Environ Med* 2009; **51**: 13–25.
- Kravchenko J, Abernethy AP, Fawzy M, Lyerly HK. Minimization of heatwave morbidity and mortality. *Am J Prev Med* 2013; **44**: 274–82.
- Xu Z, Sheffield PE, Su H, Wang X, Bi Y, Tong S. The impact of heat waves on children's health: a systematic review. *Int J Biometeorol* 2014; **58**: 239–47.
- Auger N, Fraser WD, Smargiassi A, Kosatsky T. Ambient heat and sudden infant death: a case-crossover study spanning 30 years in Montreal, Canada. *Environ Health Perspect* 2015; **123**: 712–16.
- Son JY, Lee JT, Bell ML. Is ambient temperature associated with risk of infant mortality? A multi-city study in Korea. *Environ Res* 2017; **158**: 748–52.
- Azongo DK, Awine T, Wak G, Binka FN, Oduro AR. A time series analysis of weather variability and all-cause mortality in the Kasena-Nankana Districts of Northern Ghana, 1995–2010. *Glob Health Action* 2012; **5**: 14–22.
- Diboulo E, Sié A, Rocklöv J, et al. Weather and mortality: a 10 year retrospective analysis of the Nouna Health and Demographic Surveillance System, Burkina Faso. *Glob Health Action* 2012; **5**: 6–13.
- Egondi T, Kyobutungi C, Kovats S, Muindi K, Ettarh R, Rocklöv J. Time-series analysis of weather and mortality patterns in Nairobi's informal settlements. *Glob Health Action* 2012; **5**: 23–32.
- Babalola O, Razaque A, Bishai D. Temperature extremes and infant mortality in Bangladesh: hotter months, lower mortality. *PLoS One* 2018; **13**: e0189252.
- Khalaj B, Lloyd G, Sheppard V, Dear K. The health impacts of heat waves in five regions of New South Wales, Australia: a case-only analysis. *Int Arch Occup Environ Health* 2010; **83**: 833–42.
- Lam LT. The association between climatic factors and childhood illnesses presented to hospital emergency among young children. *Int J Environ Health Res* 2007; **17**: 1–8.
- Ghirardi L, Bisoffi G, Mirandola R, Ricci G, Baccini M. The impact of heat on an emergency department in Italy: attributable visits among children, adults, and the elderly during the warm season. *PLoS One* 2015; **10**: e0141054.
- Sheffield PE, Herrera MT, Kinnee EJ, Clougherty JE. Not so little differences: variation in hot weather risk to young children in New York City. *Public Health* 2018; **161**: 119–26.
- van der Linden N, Longden T, Richards JR, et al. The use of an 'acclimatisation' heatwave measure to compare temperature-related demand for emergency services in Australia, Botswana, Netherlands, Pakistan, and USA. *PLoS One* 2019; **14**: e0214242.
- van Loenhout JAF, Delbiso TD, Kirilouk A, Rodriguez-Llanes JM, Segers J, Guha-Sapir D. Heat and emergency room admissions in the Netherlands. *BMC Public Health* 2018; **18**: 108.
- Xu Z, Huang C, Hu W, Turner LR, Su H, Tong S. Extreme temperatures and emergency department admissions for childhood asthma in Brisbane, Australia. *Occup Environ Med* 2013; **70**: 730–35.
- Otte im Kampe E, Kovats S, Hajat S. Impact of high ambient temperature on unintentional injuries in high-income countries: a narrative systematic literature review. *BMJ Open* 2016; **6**: e010399.
- Ho HC, Knudby A, Chi G, Aminipouri M, Yuk-FoLai D. Spatiotemporal analysis of regional socio-economic vulnerability change associated with heat risks in Canada. *Appl Geogr* 2018; **95**: 61–70.
- Díaz J, Carmona R, Mirón IJ, Ortiz C, Linares C. Comparison of the effects of extreme temperatures on daily mortality in Madrid (Spain), by age group: the need for a cold wave prevention plan. *Environ Res* 2015; **143**: 186–91.

- 40 Wang L, Liu T, Hu M, et al. The impact of cold spells on mortality and effect modification by cold spell characteristics. *Sci Rep* 2016; 6: 38380.
- 41 Cong X, Xu X, Zhang Y, Wang Q, Xu L, Huo X. Temperature drop and the risk of asthma: a systematic review and meta-analysis. *Environ Sci Pollut Res Int* 2017; 24: 22535–46.
- 42 Liu Y, Guo Y, Wang C, et al. Association between temperature change and outpatient visits for respiratory tract infections among children in Guangzhou, China. *Int J Environ Res Public Health* 2015; 12: 439–54.
- 43 Mills D, Jones R, Wobus C, et al. Projecting age-stratified risk of exposure to inland flooding and wildfire smoke in the United States under two climate scenarios. *Environ Health Perspect* 2018; 126: 47007.
- 44 Kabir MI, Rahman MB, Smith W, Lusha MAF, Milton AH. Climate change and health in Bangladesh: a baseline cross-sectional survey. *Glob Health Action* 2016; 9: 29609.
- 45 Mrema S, Shamte A, Selemani M, Masanja H. The influence of weather on mortality in rural Tanzania: a time-series analysis 1999–2010. *Glob Health Action* 2012; 5: 33–43.
- 46 Phung D, Huang C, Rutherford S, Chu C, Wang X, Nguyen M. Association between annual river flood pulse and paediatric hospital admissions in the Mekong Delta area. *Environ Res* 2014; 135: 212–20.
- 47 Zhang N, Song D, Zhang J, et al. The impact of the 2016 flood event in Anhui Province, China on infectious diarrhea disease: an interrupted time-series study. *Environ Int* 2019; 127: 801–09.
- 48 Rodriguez-Llanes JM, Ranjan-Dash S, Mukhopadhyay A, Guha-Sapir D. Flood-exposure is associated with higher prevalence of child undernutrition in rural eastern India. *Int J Environ Res Public Health* 2016; 13: 210.
- 49 Alpino TA, de Sena AR, de Freitas CM. Disasters related to droughts and public health—a review of the scientific literature. *Cien Saude Colet* 2016; 21: 809–20.
- 50 Dean JG, Stain HJ. Mental health impact for adolescents living with prolonged drought. *Aust J Rural Health* 2010; 18: 32–37.
- 51 Hyland M, Russ J. Water as destiny—the long-term impacts of drought in sub-Saharan Africa. *World Dev* 2019; 115: 30–45.
- 52 Bandyopadhyay S, Kanji S, Wang L, LiMin W. The impact of rainfall and temperature variation on diarrheal prevalence in sub-Saharan Africa. *Appl Geogr* 2012; 33: 63–72.
- 53 Henry SJF, Dos Santos S. Rainfall variations and child mortality in the Sahel: results from a comparative event history analysis in Burkina Faso and Mali. *Popul Environ* 2013; 34: 431–59.
- 54 Ahdoot S, Pacheco SE. Global climate change and children's health. *Pediatrics* 2015; 136: e1468–84.
- 55 Xu Z, Sheffield PE, Hu W, et al. Climate change and children's health—a call for research on what works to protect children. *Int J Environ Res Public Health* 2012; 9: 3298–316.
- 56 UNICEF. Climate change and children: a human security challenge. Florence: UNICEF, 2008.
- 57 Dasgupta S. Burden of climate change on malaria mortality. *Int J Hyg Environ Health* 2018; 221: 782–91.
- 58 Boyce R, Reyes R, Matte M, et al. Severe flooding and malaria transmission in the western Ugandan highlands: implications for disease control in an era of global climate change. *J Infect Dis* 2016; 214: 1403–10.
- 59 Ssempiira J, Kissa J, Nambuusi B, et al. Interactions between climatic changes and intervention effects on malaria spatio-temporal dynamics in Uganda. *Parasite Epidemiol Control* 2018; 3: e00070.
- 60 Semakula HM, Song G, Achuu SP, et al. Prediction of future malaria hotspots under climate change in sub-Saharan Africa. *Clim Change* 2017; 143: 415–28.
- 61 Wunderlich J, Acuña-Soto R, Alonso WJ. Dengue hospitalisations in Brazil: annual wave from west to east and recent increase among children. *Epidemiol Infect* 2018; 146: 236–45.
- 62 Akachi Y, Goodman D, Parker D. Global climate change and child health: a review of pathways, impacts and measures to improve the evidence base. Florence: UNICEF, 2009.
- 63 Ebi KL, Paulson JA. Climate change and child health in the United States. *Curr Probl Pediatr Adolesc Health Care* 2010; 40: 2–18.
- 64 Dumic I, Severnini E. 'Ticking bomb': the impact of climate change on the incidence of Lyme disease. *Can J Infect Dis Med Microbiol* 2018; 2018: 5719081.
- 65 El-Fadel M, Ghanimeh S, Maroun R, Alameddine I. Climate change and temperature rise: implications on food- and water-borne diseases. *Sci Total Environ* 2012; 437: 15–21.
- 66 Carlton EJ, Woster AP, DeWitt P, Goldstein RS, Levy K. A systematic review and meta-analysis of ambient temperature and diarrhoeal diseases. *Int J Epidemiol* 2016; 45: 117–30.
- 67 Lloyd SJ, Kovats RS, Armstrong BG. Global diarrhoea morbidity, weather and climate. *Clim Res* 2007; 34: 119–27.
- 68 Thiam S, Diène AN, Sy I, et al. Association between childhood diarrhoeal incidence and climatic factors in urban and rural settings in the health district of Mbour, Senegal. *Int J Environ Res Public Health* 2017; 14: e1049.
- 69 Uejo CK, Christenson M, Moran C, Gorelick M. Drinking-water treatment, climate change, and childhood gastrointestinal illness projections for northern Wisconsin (USA) communities drinking untreated groundwater. *Hydrogeol J* 2017; 25: 969–79.
- 70 Moors E, Singh T, Siderius C, Balakrishnan S, Mishra A. Climate change and waterborne diarrhoea in northern India: impacts and adaptation strategies. *Sci Total Environ* 2013; 468–69 (suppl): S139–51.
- 71 McIver LJ, Imai C, Buettner PG, et al. Diarrheal diseases and climate change in Cambodia. *Asia Pac J Public Health* 2016; 28: 576–85.
- 72 Thompson CN, Zeller JL, Nhu TH, et al. The impact of environmental and climatic variation on the spatiotemporal trends of hospitalized pediatric diarrhea in Ho Chi Minh City, Vietnam. *Health Place* 2015; 35: 147–54.
- 73 Phung D, Rutherford S, Chu C, et al. Temperature as a risk factor for hospitalisations among young children in the Mekong Delta area, Vietnam. *Occup Environ Med* 2015; 72: 529–35.
- 74 Mirsaeidi M, Motahari H, Taghizadeh Khamesi M, Sharifi A, Campos M, Schraufnagel DE. Climate change and respiratory infections. *Ann Am Thorac Soc* 2016; 13: 1223–30.
- 75 Xu Z, Liu Y, Ma Z, Li S, Hu W, Tong S. Impact of temperature on childhood pneumonia estimated from satellite remote sensing. *Environ Res* 2014; 132: 334–41.
- 76 Duan C, Zhang X, Jin H, et al. Meteorological factors and its association with hand, foot and mouth disease in southeast and east Asia areas: a meta-analysis. *Epidemiol Infect* 2018; 147: 1–18.
- 77 Zhao Q, Li S, Cao W, et al. Modeling the present and future incidence of pediatric hand, foot, and mouth disease associated with ambient temperature in mainland China. *Environ Health Perspect* 2018; 126: 047010.
- 78 Coates SJ, Davis MDP, Andersen LK. Temperature and humidity affect the incidence of hand, foot, and mouth disease: a systematic review of the literature—a report from the International Society of Dermatology Climate Change Committee. *Int J Dermatol* 2019; 58: 388–99.
- 79 Schweitzer MD, Calzadilla AS, Salamo O, et al. Lung health in era of climate change and dust storms. *Environ Res* 2018; 163: 36–42.
- 80 Li S, Williams G, Jalaludin B, Baker P. Panel studies of air pollution on children's lung function and respiratory symptoms: a literature review. *J Asthma* 2012; 49: 895–910.
- 81 Ward CJ. It's an ill wind: the effect of fine particulate air pollution on respiratory hospitalizations. *Can J Econ* 2015; 48: 1694–732.
- 82 Gohlke JM, Thomas R, Woodward A, et al. Estimating the global public health implications of electricity and coal consumption. *Environ Health Perspect* 2011; 119: 821–26.
- 83 Bailis R, Ezzati M, Kammen DM. Mortality and greenhouse gas impacts of biomass and petroleum energy futures in Africa. *Science* 2005; 308: 98–103.
- 84 Ha S, Zhu Y, Liu D, Sherman S, Mendola P. Ambient temperature and air quality in relation to small for gestational age and term low birthweight. *Environ Res* 2017; 155: 394–400.
- 85 Liu W, Huang C, Hu Y, et al. Associations of gestational and early life exposures to ambient air pollution with childhood respiratory diseases in Shanghai, China: a retrospective cohort study. *Environ Int* 2016; 92–93: 284–93.
- 86 Perera F, Ashrafi A, Kinney P, Mills D. Towards a fuller assessment of benefits to children's health of reducing air pollution and mitigating climate change due to fossil fuel combustion. *Environ Res* 2019; 172: 55–72.
- 87 Bernard SM, Samet JM, Grambsch A, Ebi KL, Romieu I. The potential impacts of climate variability and change on air pollution-related health effects in the United States. *Environ Health Perspect* 2001; 109 (suppl 2): 199–209.

- 88 Yoon S-J, Oh I-H, Seo H-Y, Kim E-J. Measuring the burden of disease due to climate change and developing a forecast model in South Korea. *Public Health* 2014; **128**: 725–33.
- 89 D'Amato G, Vitale C, Lanza M, Molino A, D'Amato M. Climate change, air pollution, and allergic respiratory diseases: an update. *Curr Opin Allergy Clin Immunol* 2016; **16**: 434–40.
- 90 Demain JG. Climate change and the impact on respiratory and allergic disease: 2018. *Curr Allergy Asthma Rep* 2018; **18**: 22.
- 91 D'Amato G, Holgate ST, Pawankar R, et al. Meteorological conditions, climate change, new emerging factors, and asthma and related allergic disorders. A statement of the World Allergy Organization. *World Allergy Organ J* 2015; **8**: 25.
- 92 Neumann JE, Anenberg SC, Weinberger KR, et al. Estimates of present and future asthma emergency department visits associated with exposure to oak, birch, and grass pollen in the United States. *Geohealth* 2019; **3**: 11–27.
- 93 Burke SEL, Sanson AV, Van Hoorn J. The psychological effects of climate change on children. *Curr Psychiatry Rep* 2018; **20**: 35.
- 94 Majeed H, Lee J. The impact of climate change on youth depression and mental health. *Lancet Planet Health* 2017; **1**: e94–95.
- 95 Friel S, Berry H, Dinh H, O'Brien L, Walls HL. The impact of drought on the association between food security and mental health in a nationally representative Australian sample. *BMC Public Health* 2014; **14**: 1102.
- 96 Younan D, Li L, Tuvblad C, et al. Long-term ambient temperature and externalizing behaviors in adolescents. *Am J Epidemiol* 2018; **187**: 1931–41.
- 97 Ojala M. How do children cope with global climate change? Coping strategies, engagement, and well-being. *J Environ Psychol* 2012; **32**: 225–33.
- 98 Ojala M. Coping with climate change among adolescents: implications for subjective well-being and environmental engagement. *Sustainability* 2013; **5**: 2191–209.
- 99 MacDonald JP, Willox AC, Ford JD, et al. Protective factors for mental health and well-being in a changing climate: perspectives from Inuit youth in Nunatsiavut, Labrador. *Soc Sci Med* 2015; **141**: 133–41.
- 100 Carnie T-L, Berry HL, Blinkhorn SA, Hart CR. In their own words: Young people's mental health in drought-affected rural and remote NSW. *Aust J Rural Health* 2011; **19**: 244–48.
- 101 Phalkey RK, Aranda-Jan C, Marx S, Höfle B, Sauerborn R. Systematic review of current efforts to quantify the impacts of climate change on undernutrition. *Proc Natl Acad Sci USA* 2015; **112**: E4522–29.
- 102 Swinburn BA, Kraak VI, Allender S, et al. The global syndemic of obesity, undernutrition, and climate change: the *Lancet* commission report. *Lancet* 2019; **393**: 791–846.
- 103 Thompson B, Cohen MJ, Meerman J. World food insecurity and malnutrition: scope, trends, causes and consequences. In: Thompson B, Cohen M, eds. *The impact of climate change and bioenergy on nutrition*. Dordrecht: Springer, 2012: 21–41.
- 104 Smith MR, Golden CD, Myers SS. Potential rise in iron deficiency due to future anthropogenic carbon dioxide emissions. *Geohealth* 2017; **1**: 248–57.
- 105 Datar A, Liu J, Linnemayr S, Stecher C. The impact of natural disasters on child health and investments in rural India. *Soc Sci Med* 2013; **76**: 83–91.
- 106 Alderman H. Safety nets can help address the risks to nutrition from increasing climate variability. *J Nutr* 2010; **140**: 148–52S.
- 107 Lloyd SJ, Kovats RS, Chalabi Z. Climate change, crop yields, and undernutrition: development of a model to quantify the impact of climate scenarios on child undernutrition. *Environ Health Perspect* 2011; **119**: 1817–23.
- 108 Davenport F, Grace K, Funk C, Shukla S. Child health outcomes in sub-Saharan Africa: a comparison of changes in climate and socio-economic factors. *Glob Environ Change* 2017; **46**: 72–87.
- 109 Lovo S, Veronesi M. Crop diversification and child health: empirical evidence from Tanzania. *Ecol Econ* 2019; **158**: 168–79.
- 110 Muttarak R. Too few nutrients and too many calories: climate change and the double burden of malnutrition in Asia. *Asian Popul Stud* 2019; **15**: 1–7.
- 111 Vilcins D, Sly PD, Jagals P. Environmental risk factors associated with child stunting: a systematic review of the literature. *Ann Glob Health* 2018; **84**: 551–62.
- 112 Hasegawa T, Fujimori S, Takahashi K, Yokohata T, Masui T. Economic implications of climate change impacts on human health through undernourishment. *Clim Change* 2016; **136**: 189–202.
- 113 Sheffield PE, Galvez MP. US childhood obesity and climate change: moving toward shared environmental health solutions. *Environ Justice* 2009; **2**: 207–14.
- 114 Benevolenza MA, DeRigne L. The impact of climate change and natural disasters on vulnerable populations: a systematic review of literature. *J Hum Behav Soc Environ* 2019; **29**: 266–81.
- 115 Daoud A, Halleröd B, Guha-Sapir D. Correction: what is the association between absolute child poverty, poor governance, and natural disasters? A global comparison of some of the realities of climate change. *PLoS One* 2016; **11**: e0155653.
- 116 Ahmed KJ, Haq SMA, Bartiaux F. The nexus between extreme weather events, sexual violence, and early marriage: a study of vulnerable populations in Bangladesh. *Popul Environ* 2019; **40**: 303–24.
- 117 Masson VL, Benoudji C, Reyes SS, Bernard G. How violence against women and girls undermines resilience to climate risks in Chad. *Disasters* 2019; **43** (suppl 3): S245–70.
- 118 Mathee A, Barnes B, Naidoo S, Swart A, Rother H-A. Development for children's environmental health in South Africa: past gains and future opportunities. *Dev South Afr* 2018; **35**: 283–93.
- 119 Balbus JM, Malina C. Identifying vulnerable subpopulations for climate change health effects in the United States. *J Occup Environ Med* 2009; **51**: 33–37.
- 120 Pronczuk J, Surdu S. Children's environmental health in the twenty-first century. *Ann N Y Acad Sci* 2008; **1140**: 143–54.
- 121 Randell H, Gray C. Climate change and educational attainment in the global tropics. *Proc Natl Acad Sci USA* 2019; **116**: 8840–45.
- 122 Dapi LN, Rocklöv J, Ngufacek-Tsague G, Tetanye E, Kjellstrom T. Heat impact on schoolchildren in Cameroon, Africa: potential health threat from climate change. *Glob Health Action* 2010; **3**: 5610.
- 123 Kistin EJ, Fogarty J, Pokrasso RS, McCally M, McCormick PG. Climate change, water resources and child health. *Arch Dis Child* 2010; **95**: 545–49.
- 124 Penning-Rowsell EC, Sultana P, Thompson PM. The 'last resort'? Population movement in response to climate-related hazards in Bangladesh. *Environ Sci Policy* 2013; **27**: S44–59.
- 125 Rylander C, Odland JO, Sandanger TM. Climate change and the potential effects on maternal and pregnancy outcomes: an assessment of the most vulnerable—the mother, fetus, and newborn child. *Glob Health Action* 2013; **6**: 19538.
- 126 Carolan-Olah M, Frankowska D. High environmental temperature and preterm birth: a review of the evidence. *Midwifery* 2014; **30**: 50–59.
- 127 Dadvand P, Basagaña X, Sartini C, et al. Climate extremes and the length of gestation. *Environ Health Perspect* 2011; **119**: 1449–53.
- 128 Ngo NS, Horton RM. Climate change and fetal health: the impacts of exposure to extreme temperatures in New York City. *Environ Res* 2016; **144**: 158–64.
- 129 Rocha R, Soares RR. Water scarcity and birth outcomes in the Brazilian semiarid. *J Dev Econ* 2015; **112**: 72–91.
- 130 Grace K, Davenport F, Hanson H, Funk C, Shukla S. Linking climate change and health outcomes: examining the relationship between temperature, precipitation and birth weight in Africa. *Glob Environ Change* 2015; **35**: 125–37.
- 131 Poursafa P, Keikha M, Kelishadi R. Systematic review on adverse birth outcomes of climate change. *J Res Med Sci* 2015; **20**: 397–402.
- 132 Asamoah B, Kjellstrom T, Östergren P-O. Is ambient heat exposure levels associated with miscarriage or stillbirths in hot regions? A cross-sectional study using survey data from the Ghana Maternal Health Survey 2007. *Int J Biometeorol* 2018; **62**: 319–30.
- 133 Ha S, Liu D, Zhu Y, et al. Ambient temperature and stillbirth: a multi-center retrospective cohort study. *Environ Health Perspect* 2017; **125**: 067011.
- 134 Dasgupta S, Huq M, Wheeler D. Drinking water salinity and infant mortality in coastal Bangladesh. *Water Econ Policy* 2016; **2**: 1650003.
- 135 Zeng J, Lu C, Deng Q. Prenatal exposure to diurnal temperature variation and early childhood pneumonia. *J Therm Biol* 2017; **65**: 105–12.
- 136 Miao Y, Shen Y-M, Lu C, Zeng J, Deng Q. Maternal exposure to ambient air temperature during pregnancy and early childhood pneumonia. *J Therm Biol* 2017; **69**: 288–93.

- 137 Liu W, Cai J, Huang C, et al. Associations of gestational and early life exposures to ambient air pollution with childhood atopic eczema in Shanghai, China. *Sci Total Environ* 2016; **572**: 34–42.
- 138 Nomura Y, Davey K, Pehme PM, et al. Influence of in utero exposure to maternal depression and natural disaster-related stress on infant temperament at 6 months: the children of Superstorm Sandy. *Infant Ment Health J* 2019; **40**: 204–16.
- 139 Isen A, Rossin-Slater M, Walker R. Relationship between season of birth, temperature exposure, and later life wellbeing. *Proc Natl Acad Sci USA* 2017; **114**: 13447–52.
- 140 Bunyavanich S, Landrigan CP, McMichael AJ, Epstein PR. The impact of climate change on child health. *Ambul Pediatr* 2003; **3**: 44–52.
- 141 Shea KM. Global climate change and children's health. *Pediatrics* 2007; **120**: 1149–52.
- 142 Ebi KL, Paulson JA. Climate change and children. *Pediatr Clin North Am* 2007; **54**: 213–26, vii.
- 143 Bernstein AS, Myers SS. Climate change and children's health. *Curr Opin Pediatr* 2011; **23**: 221–26.
- 144 Seal A, Vasudevan C. Climate change and child health. *Arch Dis Child* 2011; **96**: 1162–66.
- 145 Patz JA, Gibbs HK, Foley JA, Rogers JV, Smith KR. Climate change and global health: quantifying a growing ethical crisis. *EcoHealth* 2007; **4**: 397–405.
- 146 Vanos JK. Children's health and vulnerability in outdoor microclimates: a comprehensive review. *Environ Int* 2015; **76**: 1–15.
- 147 Costello A, Abbas M, Allen A, et al. Managing the health effects of climate change. *Lancet* 2009; **373**: 1693–733.
- 148 Local Burden of Disease 2017 Child Mortality Collaborators. Mapping 123 million neonatal, infant and child deaths between 2000 and 2017. *Nature* 2019; **574**: 353–58.
- 149 Alfvén T, Dahlstrand J, Humphreys D, et al. Placing children and adolescents at the centre of the Sustainable Development Goals will deliver for current and future generations. *Glob Health Action* 2019; **12**: 1670015.

Copyright © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-ND-NC 4.0 license.